

Lecture 2: Matrix Algebra Refresher

Bruno Siciliano, Lorenzo Sciavicco, Luigi Villani, and Giuseppe Oriolo. Robotics: Modelling, Planning and Control, Appendix A

CSCI 545 Introduction to Robotics
Instructor: Stefanos Nikolaidis

Matrix

- A *matrix* of dimensions ($m \times n$), with m and n positive integers, is an array of elements a_{ij} arranged into m rows and n columns

$$\mathbf{A} = [a_{ij}]_{\substack{i = 1, \dots, m \\ j = 1, \dots, n}} = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{bmatrix}$$

Matrix

- If $m = n$, the matrix is said to be *square*; if $m < n$, the matrix has more columns than rows; if $m > n$ the matrix has more rows than columns. Further, if $n = 1$, the matrix is reduced to a column vector a of dimensions $(m \times 1)$. The elements a_i are said to be vector components.

Matrix

- A square matrix A of dimensions $(n \times n)$ is said to be *upper triangular* if $a_{ij} = 0$ for $i > j$:

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ 0 & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & a_{nn} \end{bmatrix}$$

- A square matrix A of dimensions $(n \times n)$ is said to be *lower triangular* if $a_{ij} = 0$ for $i < j$:

Matrix

- An $(n \times n)$ square matrix A is said to be *diagonal* if $a_{ij} = 0$ for $i \neq j$, i.e.:

$$\mathbf{A} = \begin{bmatrix} a_{11} & 0 & \dots & 0 \\ 0 & a_{22} & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & a_{nn} \end{bmatrix} = \text{diag}\{a_{11}, a_{22}, \dots, a_{nn}\}$$

- Question: How is a matrix called if it has all unit elements in the diagonal:

Matrix

- The *transpose* A^T of a matrix A of dimensions $(m \times n)$ is the matrix of dimensions $(n \times m)$ which is obtained from the original matrix by interchanging its rows and columns.

$$A^T = \begin{bmatrix} a_{11} & a_{21} & \dots & a_{m1} \\ a_{12} & a_{22} & \dots & a_{m2} \\ \vdots & \vdots & \ddots & \vdots \\ a_{1n} & a_{2n} & \dots & a_{mn} \end{bmatrix}$$

Question

- What is the transpose of the following matrix?

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$$

$$A^T = ?$$

Matrix

- An $(n \times n)$ square matrix A is said to be *symmetric* if $A^T = A$ and thus $a_{ij} = a_{ji}$:

$$\mathbf{A} = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{12} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{1n} & a_{2n} & \dots & a_{nn} \end{bmatrix}$$

Questions

- Can you give an example of a symmetric matrix?

Questions

- Can you give an example of a symmetric matrix?

$$A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$$

Matrix

- An $(n \times n)$ square matrix A is said to be *skew-symmetric* if $A^T = -A$ and thus $a_{ij} = -a_{ji}$ for $i \neq j$ and $a_{ii} = 0$, leading to

$$A = \begin{bmatrix} 0 & a_{12} & \dots & a_{1n} \\ -a_{12} & 0 & \dots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ -a_{1n} & -a_{2n} & \dots & 0 \end{bmatrix}$$

Matrix Algebra

$C = A + B$ (for matrices that have the same shape)

$$A + B = B + A$$

$$(A + B) + C = A + (B + C)$$

Matrix Algebra

If $A_s = \frac{1}{2}(A + A^T)$ and

$A_a = \frac{1}{2}(A - A^T)$ then:

$$A = A_s + A_a$$

Example

$$A_s = \frac{1}{2}(A + A^T)$$

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \quad A^T = \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix}$$

Example

$$A_s = \frac{1}{2}(A + A^T)$$

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \quad A^T = \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix}$$

$$A_s = \begin{bmatrix} 1 & \frac{5}{2} \\ \frac{5}{2} & 4 \end{bmatrix}$$

Example

$$A_s = \frac{1}{2}(A + A^T) \quad A_a = \frac{1}{2}(A - A^T)$$

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \quad A^T = \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix}$$

$$A_s = \begin{bmatrix} 1 & \frac{5}{2} \\ \frac{5}{2} & 4 \end{bmatrix} \quad A_a = \begin{bmatrix} 0 & -\frac{1}{2} \\ \frac{1}{2} & 0 \end{bmatrix}$$

Example

$$A_s = \frac{1}{2}(A + A^T) \quad A_a = \frac{1}{2}(A - A^T)$$

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \quad A^T = \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix}$$

$$A_s = \begin{bmatrix} 1 & \frac{5}{2} \\ \frac{5}{2} & 4 \end{bmatrix} \quad A_a = \begin{bmatrix} 0 & -\frac{1}{2} \\ \frac{1}{2} & 0 \end{bmatrix}$$

$$A = A_s + A_a$$

Multiplication of Matrices

$$A(BC) = (AB)C$$

$$A(B+C) = AB + AC$$

$$(A+B)C = AC + BC$$

$$(AB)^T = B^T A^T$$

Multiplication of Matrices

$$*AB \neq BA*$$

Example

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \quad B = \begin{bmatrix} 2 & 2 \\ 2 & 3 \end{bmatrix}$$

Example

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \quad B = \begin{bmatrix} 2 & 2 \\ 2 & 3 \end{bmatrix}$$

$$AB = \begin{bmatrix} 6 & 8 \\ 14 & 18 \end{bmatrix}$$

Example

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \quad B = \begin{bmatrix} 2 & 2 \\ 2 & 3 \end{bmatrix}$$

$$AB = \begin{bmatrix} 6 & 8 \\ 14 & 18 \end{bmatrix}$$

$$BA = \begin{bmatrix} 8 & 12 \\ 11 & 16 \end{bmatrix}$$

Multiplication of Matrices


$$AB \neq BA$$

Linear Independence

- Rank: This corresponds to the maximal number of linearly independent columns of A .
- A set of vectors is said to be **linearly dependent** if at least one of the vectors in the set can be defined as a linear combination of the others.
- If no vector in the set can be written in this way, then the vectors are said to be **linearly independent**.

Interpretation of Matrices

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$




What is the rank of this matrix?

Interpretation of Matrices

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\det(A) = 0$$

What is the rank of this matrix?



Inverse Matrices

$$A^{-1}A = I$$

Matrix Transformations

Vector \mathbf{w} is transformed to a new space

$$A\mathbf{w} = \mathbf{v}$$

Vector \mathbf{w} is transformed back to the old space

$$\mathbf{w} = A^{-1}\mathbf{v}$$

But what if \mathbf{A} is rank deficient?

Orthogonal Matrices

If A is a square matrix, and:

$$A^T = A^{-1}$$

then the matrix is called **orthogonal**.

Properties of Inverse Matrices

If A and B are square invertible matrices:

$$(AB)^{-1} = B^{-1}A^{-1}$$

Matrix Derivatives

The *derivative* of an $(m \times n)$ matrix $A(t)$, whose elements $a_{ij}(t)$ are differentiable functions, is the matrix

$$\dot{A}(t) = \frac{d}{dt}A(t) = \left[\frac{d}{dt}a_{ij}(t) \right]_{i=1, \dots, m, j=1, \dots, n}$$

Example

$$A = \begin{bmatrix} t & t^2 \\ 7t & 1 \end{bmatrix} \dot{A}(t) = ?$$

Example

$$\dot{A}(t) = \begin{bmatrix} 1 & 2t \\ 7 & 0 \end{bmatrix}$$

Gradient of a Function

$$\nabla_x f(x) = \left(\frac{\partial f(x)}{\partial x} \right) = \left[\frac{\partial f(x)}{\partial x_1} \quad \frac{\partial f(x)}{\partial x_2} \quad \dots \quad \frac{\partial f(x)}{\partial x_n} \right]$$

Example

$$f(x_1, x_2) = x_1 * x_2 + x_1^2$$

$$\nabla_x f(x) = ?$$

Example

$$f(x_1, x_2) = x_1 * x_2 + x_1^2$$

$$\begin{aligned}\nabla_x f(x) &= \left[\frac{\partial f(x)}{\partial x_1}, \frac{\partial f(x)}{\partial x_2} \right] \\ &= [x_2 + 2x_1, x_1]\end{aligned}$$

Gradient of a Function

$$\dot{f}(x) = \frac{d}{dt} f(x) = \frac{\partial f}{\partial x} \dot{x} = \nabla_x f(x) \dot{x}$$

Example

$$f(x_1, x_2) = x_1^2 + x_2^2$$

$$x_1(t) = 3t$$

$$x_2(t) = 4t^2$$

Example

$$f(x_1, x_2) = x_1^2 + x_2^2$$

$$x_1(t) = 3t$$

$$x_2(t) = 4t^2$$

$$\dot{f}(x) = \frac{d}{dt} f(x)$$

Example

$$f(x_1, x_2) = x_1^2 + x_2^2$$

$$x_1(t) = 3t$$

$$x_2(t) = 4t^2$$

$$\dot{f}(x) = \frac{d}{dt} f(x)$$

$$= \frac{\partial f}{\partial x} \dot{x}$$

$$= \nabla_x f(x) \dot{x}$$

$$= (2x_1, 2x_2)^T (3, 8t)$$

$$= (6t, 8t^2)^T (3, 8t)$$

$$= 18t + 64t^3$$

Jacobian Matrix

$$J_g(x) = \frac{g(x)}{\partial x} = \begin{bmatrix} \frac{\partial g_1(x)}{\partial x} \\ \frac{\partial g_2(x)}{\partial x} \\ \vdots \\ \frac{\partial g_m(x)}{\partial x} \end{bmatrix}$$

Example

$$g = [g_1, g_2]$$

$$g_1(x_1, x_2) = x_1 * x_2 + x_1^2$$

$$g_2(x_1, x_2) = x_1^4$$

Example

$$g = [g_1, g_2]$$

$$g_1(x_1, x_2) = x_1 * x_2 + x_1^2$$

$$g_2(x_1, x_2) = x_1^4$$

$$\frac{\partial g_1}{\partial x} = [x_2 + 2x_1, x_1]$$

$$\frac{\partial g_2}{\partial x} = [4 * x_1^3, 0]$$

Example

$$g = [g_1, g_2]$$

$$g_1(x_1, x_2) = x_1 * x_2 + x_1^2$$

$$g_2(x_1, x_2) = x_1^4$$

$$\frac{\partial g_1}{\partial x} = [x_2 + 2x_1, x_1]$$

$$\frac{\partial g_2}{\partial x} = [4 * x_1^3, 0]$$

$$J = \begin{bmatrix} x_2 + 2x_1 & x_1 \\ 4x_1^3 & 0 \end{bmatrix}$$

Time Derivative of Vector Function

$$\dot{g}(x) = \frac{d}{dt}g(x(t)) = \frac{\partial g}{\partial x}\dot{x} = J_g(x)\dot{x}$$


Inner Product

Given n vectors x_i of dimensions $(m \times 1)$, the **inner product** is defined as :

$$\langle x, y \rangle = x_1y_1 + x_2y_2 + \dots + x_my_m = x^T y = y^T x$$

Inner Product: Geometric Interpretation

$$x^T y = \|x\| \|y\| \cos(\theta)$$



What is the geometric interpretation of the dot product if the vector y has length = 1?

Orthogonal vectors


- Two vectors are said to be *orthogonal* when their scalar product is 0:

$$x^T y = 0$$

Cross Product

$$x \times y = \begin{bmatrix} x_2 y_3 - x_3 y_2 \\ x_3 y_1 - x_1 y_3 \\ x_1 y_2 - x_2 y_1 \end{bmatrix}$$

Cross Product



Question: How do you get the cross-product of two vectors to be 0?

Cross Product Operator

$$S(x) = \begin{bmatrix} 0 & -x_3 & x_2 \\ x_3 & 0 & -x_1 \\ -x_2 & x_1 & 0 \end{bmatrix}$$

Cross Product Operator

$$x \times y = S(x)y = \begin{bmatrix} 0 & -x_3 & x_2 \\ x_3 & 0 & -x_1 \\ -x_2 & x_1 & 0 \end{bmatrix} y$$

Eigenvalues and Eigenvectors

- There are some vector u , that if you multiply with A they either stretch or become shorter, but they stay in the same direction of motion.

$$Au - \lambda u = 0$$

Eigenvalues and Eigenvectors

- There are some vector u , that if you multiply with A they either stretch or become shorter, but they stay in the same direction of motion.

$$Au - \lambda u = 0$$

$$(A - \lambda I)u = 0$$

Eigenvalues and Eigenvectors

- There are some vector u , that if you multiply with A they either stretch or become shorter, but they stay in the same direction of motion.

$$Au - \lambda u = 0$$

$$(A - \lambda I)u = 0$$

$$\det(A - \lambda I) = 0$$

Example

$$A = \begin{bmatrix} 0 & 1 \\ -2 & -3 \end{bmatrix}$$

Example

$$A = \begin{bmatrix} 0 & 1 \\ -2 & -3 \end{bmatrix}$$

$$\det(\lambda I - A) = \det\left(\begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix} - \begin{bmatrix} 0 & 1 \\ -2 & -3 \end{bmatrix}\right)$$

Example

$$A = \begin{bmatrix} 0 & 1 \\ -2 & -3 \end{bmatrix}$$

$$\det(\lambda I - A) = \det\left(\begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix} - \begin{bmatrix} 0 & 1 \\ -2 & -3 \end{bmatrix}\right)$$

$$= \det\left(\begin{bmatrix} \lambda & -1 \\ 2 & \lambda + 3 \end{bmatrix}\right)$$

$$= \lambda^2 + 3\lambda + 2 = 0$$

Example

$$A = \begin{bmatrix} 0 & 1 \\ -2 & -3 \end{bmatrix}$$

$$\det(\lambda I - A) = \det\left(\begin{bmatrix} \lambda & 0 \\ 0 & \lambda \end{bmatrix} - \begin{bmatrix} 0 & 1 \\ -2 & -3 \end{bmatrix}\right)$$

$$= \det\left(\begin{bmatrix} \lambda & -1 \\ 2 & \lambda + 3 \end{bmatrix}\right)$$

$$= \lambda^2 + 3\lambda + 2 = 0$$

$$\lambda_1 = -2, \lambda_2 = -1$$

Positive Definiteness

- The matrix A is positive definite iff:

$$x^T A x > 0, \forall x \neq 0$$

$$x^T A x = 0, \forall x = 0$$

Example

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$$

Example

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$$

$$\begin{aligned} [x_1, x_2] \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix} [x_1, x_2]^T &= [x_1, 2x_2][x_1, x_2]^T \\ &= x_1^2 + 2x_2^2 \end{aligned}$$